



tradecorp Ca

Premium Calcium (Ca) EDTA Chelate as 100% soluble micro-granule to prevent & correct Calcium deficiency



DESCRIPTION

- Calcium (Ca) EDTA Chelate with maximum Chelation %*
- 100% soluble micro-granule (WG) for rapid & instantaneous solubility
- Corrects and prevents Calcium deficiencies in all crops
- Use on all crops from pre-flowering to harvest for quality fruit & vegetables
- Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
- Suitable for use in soils & solutions with pH 4-9
- Production system certified to ISO 9001 & ISO 14001
- Guaranteed Contents
- Ecocert Approved
- 5 kg & 1 kg bag size


(* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

TECHNICAL SPECIFICATIONS

- Calcium (Ca) Chelated with EDTA and soluble in water: 10%
- Chelating agent: EDTA
- Presentation: Soluble micro-granules (WG)
- pH stability range: 4-9
- Solubility: 300 g /L

Tradecorp APAC Pty. Ltd.
Level 1, 225 George Street, Sydney - NSW 2000
Contact: 0448 016 025
Email: australia@tradecorp.sapec.pt





MAIN BENEFITS

- Single element EDTA chelates allow for precise and independent nutrient adjustment & dosage
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Easy and convenient handling without lumps or sediments
- Convenient micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

FOLIAR APPLICATION

General dosage	0.5-1.5 kg /ha	2-4 applications before or as soon as symptoms appear 14 days apart
Vegetables (e.g. tomato, cucurbits)	0.5-1.5 kg /ha (75-100 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Tubers (e.g. Potato)	1 kg /ha	3-4 applications every 7-10 days once stems reach 10 cm. Use high volumes of water (e.g. 500 L)
Orchard Crops	0.5-1.5 kg /ha (50-100 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Cereals	0.5-1 kg /ha	2-3 applications at tillering, stem elongation and flag leaf
Strawberry	0.5-1 kg /ha	2-6 applications every 7-10 days once foliage is sufficient
Ornamentals	0.25-0.5 kg /ha	2-3 applications every 7-14 days once symptoms appear

DOSAGE AND METHOD OF USE

DRIP APPLICATION

General Dose	1-2.5 kg /ha	2-6 applications early in the crop cycle every 14 days
---------------------	--------------	--

HYDROPONICS

General Dose	0.4-0.6 kg /1000 L (x100)	Use 1 L of mother stock solution /100 L irrigation water
---------------------	------------------------------	--

SOIL APPLICATION

General Dose	5-10 kg /ha	1 application before sowing, transplant or the beginning of plant growth
---------------------	-------------	--