



tradecorp Fe

Premium Iron EDTA chelate as 100% soluble wettable powder to prevent & correct Iron deficiency



DESCRIPTION

- Iron (Fe) EDTA Chelate with maximum Chelation %*
- 100% soluble wettable powder (WP) for rapid & instantaneous solubility
- Corrects and prevents Iron deficiency in all crops
- Most Iron should be applied by fruit set
- Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
- Use on all crops by foliar, and in soils with pH 4-7
- Production system certified to ISO 9001 & ISO 14001
- Guaranteed Contents
- Ecocert Approved
- 5 kg & 1 kg bag size


(* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

TECHNICAL SPECIFICATIONS

- | | |
|--|---------------------|
| - Iron (Fe) chelated with EDTA and soluble in water: | 13.2% w/w |
| - Chelating agent: | EDTA |
| - Presentation: | Soluble powder (WP) |
| - pH stability range: | 4-7 |
| - Solubility: | 70 g /L |

Tradecorp APAC Pty. Ltd.
Level 1, 225 George Street, Sydney - NSW 2000
Contact: 0448 016 025
Email: australia@tradecorp.sapec.pt





MAIN BENEFITS

- Single element EDTA Chelates enable precise and independent nutrient adjustment & dosage
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring wettable powder (WP), no rinsing of packing
- High efficiency at low dose rates

FOLIAR APPLICATION

General dose	0.5-1.5 kg /ha	2-3 applications before or as soon as symptoms appear 14 days apart
Vegetables (e.g. tomato, cucurbits)	1-1.5 kg /ha (75-100 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Tubers (e.g. Potato)	1-1.5 kg /ha	2-3 applications once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500 L)
Orchard Crops	0.5-1 kg /ha (30-100 g/100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Vines	1-1.5 kg /ha	3-4 applications 2 applications before flower and 2 after flowering
Cereals & Herbaceous	1 kg /ha	2-3 applications at tillering, stem elongation and flag leaf
Ornamentals	10-25 g /100 L	2-4 applications every 7-10 days from beginning of vegetative cycle

DOSAGE AND METHOD OF USE

DRIP APPLICATION

General Dose	2-4 kg /ha	2-4 applications early in the crop cycle up until fruit fill 14 days apart
Vines	3-5 kg /ha	3-4 applications from budburst

HYDROPONICS

General Dose	0.4-0.6 kg /1000 L (x100)	Use 1 L of mother stock solution /100 L irrigation water
---------------------	------------------------------	--

SOIL APPLICATION

General Dose	4-8 kg /ha	1 application before sowing, transplant or the beginning of plant growth
---------------------	------------	--