



ultraferro

The premier Iron (Fe) EDDHA formulation to prevent & correct Iron Chlorosis in All Soils, 100% water soluble



DESCRIPTION

- Iron (Fe) EDDHA Chelate with maximum Chelation %*
- 100% soluble micro-granule (WG) for rapid & instantaneous solubility
- Corrects and prevents Iron deficiencies in All Soils
- Use on all crops, the majority of Iron should be applied before flowering
- Suitable for use in drip, direct soil, substrate & hydroponic applications
- Suitable for use in soils & solutions with pH 3-11
- Production system certified to ISO 9001 & ISO 14001
- Guaranteed Contents
- Ecocert Approved
- 5 kg & 1 kg bag size

(* Chelation % is the % of the micronutrient attached to the EDDHA, higher is better)

TECHNICAL SPECIFICATIONS

- | | |
|--|-----------------------------|
| - Iron (Fe) chelated with EDDHA and soluble in water: | 6% w/w |
| - Iron (Fe) chelated with EDDHA in ortho-ortho position: | 4.2% w/w |
| - Chelating agent: | EDDHA |
| - Presentation: | Soluble micro-granules (WG) |
| - pH stability range: | 3-11 |
| - Solubility: | 60 g /L |

Tradecorp APAC Pty. Ltd.
Level 1, 225 George Street, Sydney - NSW 2000
Contact: 0448 016 025
Email: australia@tradecorp.sapec.pt





MAIN BENEFITS

- Single element Iron EDDHA Chelates enable precise and independent nutrient adjustment & dosage
- Iron EDDHA Chelates are the most efficient form of Iron nutrient for soil use
- EDDHA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

FERTIGATION

| | | |
|---------------------|----------|--|
| General Dose | 2 kg /ha | 2-6 applications split from early vegetative stage until fruit set |
|---------------------|----------|--|

HYDROPONICS

| | | |
|---------------------|------------------------|---|
| General Dose | 1-1.5kg /1000 L (x100) | 2-4 applications early in the crop cycle up until fruit fill. 14 days apart |
|---------------------|------------------------|---|

SOIL APPLICATION

| | | |
|-----------------------------------|-----------------------|--|
| General dose | 5-10 kg /ha | 1 application as a single soil spray or soil injection early in the season where fertigation is not possible |
| Orchard (Establishment) | 5 g /tree | 2 applications at establishment |
| Orchard (Early Production) | 10-20 g /tree | 2 applications pre-flowering |
| Orchard (Full Production) | 25-50 g /tree | |
| Vegetables | 1-7g /m ² | 1-3 applications before sowing, transplant or the beginning of plant growth |
| Flower Beds | 3-6 g /m ² | 1-3 applications early in the season as required |

FLOWERS

| | | |
|---------------------|----------|--------|
| General Dose | 0.5 g /L | Weekly |
|---------------------|----------|--------|

DOSAGE AND METHOD OF USE