



tradecorp AZ fresco

Premium chemical mix of 6 micro-nutrient EDTA Chelates with Magnesium & Sulfur as 100% soluble micro-granule to prevent & correct multiple deficiencies



DESCRIPTION

- Premium 6 micro-nutrient EDTA chemical mix with emphasis on Iron (Fe) 4%, Zinc (Zn) 4% and Manganese (Mn) 3%
 - Corrects and prevents micro-nutrient deficiencies in all crops
 - EDTA Chelates with maximum Chelation %*
 - 100% soluble micro-granule (WG) for rapid & instantaneous solubility
 - Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
 - Suitable for use in soils & solutions with pH 3-9
 - Production system certified to ISO 9001 & ISO 14001
 - Guaranteed Contents, Ecocert Approved
 - 5 kg bag size
- (* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

TECHNICAL SPECIFICATIONS

- | | |
|---|-----------|
| - Magnesium (Mg) soluble in water: | 1.20% w/w |
| - Sulfur (S) soluble in water: | 2.40% w/w |
| - Zinc (Zn) soluble in water and chelated by EDTA: | 4.0% w/w |
| - Iron (Fe) soluble in water and chelated by EDTA: | 4.0% w/w |
| - Manganese (Mn) soluble in water and chelated by EDTA: | 3.0% w/w |
| - Copper (Cu) soluble in water and chelated by EDTA: | 0.60% w/w |
| - Boron (B) soluble in water: | 1.50% w/w |
| - Molybdenum (Mo) soluble in water: | 0.05% w/w |
| - EDTA pH stability range: | 3-9 |

Tradecorp APAC Pty. Ltd.
 Level 1, 225 George Street, Sydney - NSW 2000
 Contact: 0448 016 025
 Email: australia@tradecorp.sapec.pt

 **tradecorp**
 nutri-performance



tradecorp AZ fresco



MAIN BENEFITS

- Chemical premix that tackles all micronutrient deficiencies
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

FOLIAR APPLICATION

General Dose	0.5-1.5 kg /ha	2-3 applications before or as soon as symptoms appear 14 days apart
Vegetables (e.g. tomato, cucurbits)	0.5-1 kg /ha (75-100 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Tubers (e.g. Potato)	1 kg /ha	3-4 applications once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500 L)
Orchard Crops	1-1.5 kg /ha	2-4 applications before flowering x2-3 and during fruit set x1
Cereals	0.5-1 kg /ha	2-3 applications at start of tillering, stem elongation and flag leaf
Ornamentals	0.2-0.5 kg /ha (75-100 g /100 L)	2-3 applications every 7-10 days from beginning of vegetative cycle

DOSAGE AND METHOD OF USE

FERTIGATION

General Dose	2-3 kg /ha	2-4 applications every 7-10 days from early in the crop cycle
---------------------	------------	---

HYDROPONICS

General Dose	3-5 kg /1000 L (x100)	Use 1 L of mother stock solution /100 L irrigation water. Use the higher rate from fruit set, high temperatures or shorter winter days
---------------------	-----------------------	--

SOIL APPLICATION

General Dose	5-10 kg /ha	1 application before sowing, transplant or the beginning of plant growth
---------------------	-------------	--