


NUTRITIONAL PROGRAM


MANGO



 soil application
 foliar application









OBJECTIVE	Post Harvest	Leaf Flush Folded	Leaf Flush Hardening	Flowering	Petal Fall Fruit Set	Fruit Swell	Ripening	Harvest
-----------	--------------	-------------------	----------------------	-----------	----------------------	-------------	----------	---------

Root growth	 humifirst WG 2 kg /ha 3-4 applications every 21 days							
-------------	---	--	--	--	--	--	--	--

Vegetative growth	 intake 2 L /ha 2-4 applications every 14 days							
-------------------	--	--	--	--	--	--	--	--

Vegetative strength	 trafos synergy 3 L /ha 1 application	 trafos synergy	3 L /ha 3-4 applications every 21 days					
---------------------	---	--	--	--	--	--	--	--

Cell division Pollen viability	  tradecorp Zn 0.1% as foliar or 2 kg /ha	  tradecorp Zn	0.1% as foliar or 2 kg /ha soil application					
-----------------------------------	--	--	---	--	--	--	--	--

Prevent micronutrient Deficiencies	 tradecorp AZ plus 2 kg /ha 2 applications 14-21 days apart		 tradecorp AZ plus 1-2 kg /ha 2 applications 14-21 days apart					
------------------------------------	---	--	---	--	--	--	--	--

Pollen germination Prevent crackings	 tradebor 2 L /ha 1 application		 tradebor	2-3 L /ha 3 applications every 14-21 days				
---	---	--	--	---	--	--	--	--

Reduce fruit drop			 systemag 1.5-2 kg /ha 1-2 applications before flowering, during petal fall and fruit set					
-------------------	--	--	--	--	--	--	--	--

Vigorous flowering			 boramin Ca 3-4 L /ha 2-3 applications during and post flowering					
--------------------	--	--	--	--	--	--	--	--

Vigorous flowering Abiotic stress			 delfan plus 3 L /ha 2-4 applications every 14 days					
--------------------------------------	--	--	---	--	--	--	--	--

Fruit quality			 systacal plus	2 kg /ha 3-5 applications every 14-21 days				
---------------	--	--	---	--	--	--	--	--

Fruit fill & quality			 amifol K	3-5 L /ha 3-5 applications every 14-21 days				
----------------------	--	--	--	---	--	--	--	--



Use Lower 7 to acidify and condition the irrigation water
 Adjust foliar spray solution to a pH of 5.8 and neutralise bicarbonates with Spray Plus for maximum foliar application efficiency

www.tradecorpastralia.com.au

