



tradecorp Zn

Premium Zinc (Zn) EDTA Chelate as 100% soluble micro-granule to prevent & correct Zinc deficiency



DESCRIPTION

- Zinc (Zn) EDTA Chelate with maximum Chelation %*
- 100% soluble micro-granule (WG) for rapid & instantaneous solubility
- Corrects and prevents Zinc deficiencies in all crops
- Use on all crops, the majority of Zinc should be applied before flowering
- Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
- Suitable for use in soils & solutions with pH 4-9
- Production system certified to ISO 9001 & ISO 14001
- Guaranteed Contents
- Ecocert Approved
- 5 kg & 1 kg bag size

(* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

TECHNICAL SPECIFICATIONS

- Zinc (Zn) chelated with EDTA and soluble in water: 14% w/w
- Chelating agent: EDTA
- Presentation: Soluble micro-granules (WG)
- pH stability range: 4-9
- Solubility: 700 g/L


Tradecorp APAC Pty. Ltd.

- U11 /20 Jijaws Street, Sumner, Brisbane - QLD 4074
 - Level 1, 225 George Street, Sydney - NSW 2000
 Tel: 1300 595 000 australia@tradecorp.sapec.pt





tradecorp Zn



MAIN BENEFITS

- Single element EDTA Chelates enable precise and independent nutrient adjustment & dosage
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

FOLIAR APPLICATION

General Dosage	0.5-1.5 kg /ha	2-3 applications before or as soon as symptoms appear 14 days apart
Vegetables (e.g. tomato, cucurbits)	1-1.5 kg /ha (75-100 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient. The majority of Zn applications should be completed before flowering
Tubers (e.g. Potato)	1-1.5 kg /ha	3-4 applications once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500 L)
Orchard Crops	1-1.5 kg /ha	2-3 applications early in the crop cycle once foliage is sufficient. The majority of Zn applications should be completed before flowering Apply 1-2 applications immediately post harvest for next years flower and pollen development
Cereals & Herbaceous	0.5-1 kg /ha	2-3 applications at tillering, stem elongation and flag leaf
Ornamentals	75-100 g /100 L	2-3 applications every 7-10 days from beginning of vegetative cycle

DOSAGE AND METHOD OF USE

DRIP APPLICATION

General Dosage	2-4 kg /ha	2-4 applications early in the crop cycle up until fruit fill. 14 days apart
-----------------------	------------	---

HYDROPONICS

General Dosage	0.3-0.5 kg /1000 L (x100)	Use 1 L of mother stock solution /100 L irrigation water
-----------------------	------------------------------	--

SOIL APPLICATION

General Dosage	4-8 kg /ha	1 application before sowing, transplant or the beginning of plant growth
-----------------------	------------	--