



tradecorp Cu

Premium Copper (Cu) EDTA Chelate as 100% soluble micro-granule to prevent & correct Copper deficiency



DESCRIPTION

- Copper (Cu) EDTA Chelate with maximum Chelation %*
 - 100% soluble micro-granule (WG) for rapid & instantaneous solubility
 - Corrects and prevents Copper deficiencies in all crops
 - Use on all crops, especially crops where seed formation is important e.g. cereals
 - Suitable for use in foliar, drip, direct soil, substrate & hydroponic applications
 - Suitable for use in soils & solutions with pH 4-9
 - Production system certified to ISO 9001 & ISO 14001
 - Guaranteed Contents
 - Ecocert Approved
 - 5 kg & 1 kg bag size
- (* Chelation % is the % of the micronutrient attached to the EDTA, higher is better)

TECHNICAL SPECIFICATIONS

- Copper (Cu) chelated with EDTA and soluble in water: 14.5% w/w
- Chelating agent: EDTA
- Presentation: Soluble micro-granules (WG)
- pH stability range: 4-9
- Solubility: 700 g /L

Tradecorp APAC Pty. Ltd.
 Level 1, 225 George Street, Sydney - NSW 2000
 Contact: 0448 016 025
 Email: australia@tradecorp.sapec.pt





MAIN BENEFITS

- Single element EDTA Chelates enable precise and independent nutrient adjustment & dosage
- EDTA Chelates are the most efficient form of nutrient for soil and foliar use
- EDTA Chelates remain soluble in the soil solution for continued availability
- Compatible with most commonly used fertilizers and agrochemical products
- Simple and convenient handling without lumps or sediments
- Easy pouring micro-granule (WG), no rinsing of packing
- High efficiency at low dose rates

FOLIAR APPLICATION

General dosage	0.5-0.75 kg /ha	2-3 applications before or as soon as symptoms appear. 14 days apart
Vegetables (e.g. tomato, cucurbits)	0.5-0.75 kg /ha (15-40 g /100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Tubers (e.g. Potato)	0.5-1 kg /ha	3-4 applications once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500 L)
Orchard Crops	0.75-1.0 kg /ha (90-100 g/100 L)	2-3 applications early in the crop cycle once foliage is sufficient
Cereals & Herbaceous	0.75-1.25 kg /ha	2 applications at tillering & stem elongation. Min. 200 L /ha water
Ornamentals	10-40 g /100 L	2-4 applications every 7-10 days from beginning of vegetative cycle appear

DOSAGE AND METHOD OF USE

DRIP APPLICATION

General Dose	0.75-1.5 kg /ha	1-3 applications early in the crop cycle up until fruit fill 14 days apart
---------------------	-----------------	--

HYDROPONICS

General Dose	0.3-0.6 kg /1000 L (x100)	Use 1 L of mother stock solution /100 L irrigation water
---------------------	------------------------------	--

SOIL APPLICATION

General Dose	1-4 kg /ha	1 application before sowing, transplant or the beginning of plant growth. Use the lower rate when applied during sowing
---------------------	------------	---