# tradecorp<sup>®</sup> Ca

Premium Calcium (Ca) EDTA chelate as 100% soluble micro-granule to prevent & correct Calcium deficiency





# **GUARANTEED CONTENTS:**

% w/w

Calcium (Ca) soluble in water and chelated by EDTA:

10

pH stability range: 4-9

Tradecorp APAC Pty. Ltd. Level 1, 225 George Street Sydney - NSW 2000 Contact: 0448 016 025

Email: australia@tradecorp.sapec.pt www.tradecorpaustralia.com.au

Made in Spain by: Trade Corporation International S.A.U. C/ Alcalá, 498, 2ª Planta 28027 Madrid (España) Tel.: (+34) 913 273 200 www.tradecorp.com.es e-mail: global@tradecorp.com.es

# tradecorp Ca

## DOSAGE AND METHOD OF USE

Application	Crop	Rate (kg /Ha)	No. of Applications	Comments
Foliar	General Dose	0.5 - 1.5	2 - 4	Apply before or as soon as symptoms appear. 14 days apart.
	Vegetables (tomato, cucurbits)	0.5 - 1.5 (75 - 100g /100L)	2 - 3	Apply early in the crop cycle once foliage is sufficient.
	Tubers (e.g. Potato)	1	3 - 4	Apply every 7 - 10 days once stems reach 10 cm. Use high volumes of water (e.g. 500L).
	Orchard Crops	0.5 - 1.5 (50 - 100g /100L)	2 - 3	Apply early in the crop cycle once foliage is sufficient.
	Cereals & Herbaceous	0.5 - 1	2 - 3	Apply at start of tillering, stem elongation and flag leaf.
	Strawberry	0.5 - 1	2 - 6	Every 7 - 10 days once foliage is sufficient.
	Ornamentals	0.25 - 0.5	2 - 3	Every 7 - 14 days once symptoms appear.
Drip	General Dose	1 - 2.5	2 - 6	Apply early in the crop cycle every 14 days.
Hydroponics	General Dose	0.4 - 0.6 kg /1000L (x100)	-	Use 1L of mother stock solution /100L irrigation water.
Soil	General Dose	5 - 10	1	Apply before sowing, transplant or the beginning of plant growth.

Spray onto dry foliage for maximum efficiency

### **DIRECTIONS FOR USE**

Do not apply foliar EDTA chelates to fruit crops at >0.1% concentration (100 g /100L water). This is cumulative. If you mix two different EDTA chelate products you must not exceed 100 g /100L for the two products. i.e. maximum dosage will be 50 g of each product to give combined 100g /100L water.

Do not apply foliar EDTA chelates to broad-acre crops at >0.5% concentration (0.5 kg /100L water). The same warning about cumulative dosage applies when mixing two different EDTA products as for fruit crops above.

Do not apply by foliar application to plum or apricot trees.

Do not apply during flowering or during fruit colour change.

Add Tradecorp Ca to water while agitating and continue agitation until granules have fully dissolved.

### **COMPATIBILITY**

Tradecorp Ca is compatible with most commonly used fertilizers and agrochemicals. It is recommended to carry out a test before preparing any mixtures and to consult technically authorized personnel.

### WARNINGS, HANDLING AND STORAGE

Keep out of reach of children. Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures. Keep only in original container. Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians.

Disclaimer: The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.



Manufacturing date: Lot number:

EN18/2V02/1016