

tradebor

Boron ethanamine preventive & remedial Boron deficiency corrector



EC FERTILIZER

Guaranteed contents:

| | % w/v |
|------------|-------|
| Boron (B): | 15.4 |

EN182/02/0315

Distributor:

Tradecorp APAC Pty. Ltd.
Level 1, 225 George Street
Sydney – NSW 2000

Contact: 0448 016 025

Email: australia@tradecorp.sapec.pt

www.tradecorpastralia.com.au

NET CONTENT
20 L (28 Kg)



COMPANY WITH
MANAGEMENT SYSTEM
CERTIFIED BY DNV
= ISO 9001 =
= ISO 14001 =



Tradebor is a concentrated specialty Boron (15.4% w/v) fertilizer complexed with ethanolamine and free from Sodium typically found in Boron fertilizers. The ethanolamine complex enhances absorption through the leaf and translocation inside the plant. As such Tradebor is particularly recommended as a foliar fertilizer although it can also be applied by drip. The following crops are particularly demanding for Boron: Almond, Avocado, Brassica, Cucurbits, Grape, Mango, Potato, Pome Fruit, Salad leaves. The advantages of Tradebor are:

- Enhanced uptake and increased translocation compared to other Boron formulations
- Easy OHS compliance, Tradebor does not have a Category 1B classification under GHS
- No sodium commonly found in other Boron fertilizers

DOSAGE AND METHOD OF USE

| Application | Crop | Rate (L /Ha) | No. of Applications | Comments |
|-------------|---------------------------|--------------|---------------------|---|
| Foliar | General (Horticulture) | 1 - 2 | 2 | Apply when the crop has sufficient vegetation, and before flowering. Repeat if necessary with minimum 10 - 14 day interval. |
| | General (Orchard & Vines) | 1 - 3 | 2 - 4 | Apply when buds are opening, before flowering, during fruit set and early fruit fill. Always apply 1 - 2 applications post harvest. Minimum 14 day interval between sprays. |
| | Strawberry | 1 | 2 - 4 | Apply at white bud stage and early flowerings. |
| | Cucurbits (e.g. Melons) | 1 - 2 | 2 - 4 | Apply when crop is 10 - 15 cm (4 - 6 leaf) before flowering and at fruit set. |
| | Potato | 2 - 4 | 2 | Begin applications at row closure and continue during early tuber formation |
| | Brassica & Salad Crops | 1 - 2 | 2 - 3 | Apply from 4 true leaves 10 - 14 days apart |
| | Grape | 2 - 3 | 1 - 3 | Apply when flowers are visible, fruit set and at veraison. |
| | Almond and Nuts | 1 - 2 | 2 - 3 | Apply during nut set and development. Always apply 1 - 2 applications post harvest for bud development. |
| | Legumes | 1 - 2 | 1 - 2 | Apply when the crop reaches 10 - 15 cm tall |
| | Lucerne | 1 - 3 | 2 - 4 | Apply when the crop reaches 10 - 15 cm tall and again 14 days later. Thereafter at every alternate cut |
| Drip | General | 2 - 4 | 2 - 4 | Apply as required |

Spray onto dry foliage using a minimum of 150 – 200 L /Ha to ensure optimum coverage of the leaf and foresee 3 – 4 hours without rain for optimal penetration.

COMPATIBILITY

Tradebor is compatible with most commonly used fertilizers and agrochemical products. It is recommended to carry out a test before preparing any mixture and to consult technically authorized personnel.

WARNINGS, HANDLING AND STORAGE

SHAKE BEFORE USE. Keep out of reach of children. Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures. Keep only in original container. Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

Disclaimer: The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.

Manufacturing date:
Lot number:



EN182/02/0315

Suitable product for Organic Agriculture according to EC Regulation No. 834/2007 and 889/2008 and to NOP rules. Control ECOCERT SA F-32600