

# tradecorp<sup>®</sup> AZ fresco

Premium chemical mix of 6 micro nutrient EDTA chelates with Magnesium & Sulfur as 100% soluble micro-granule to prevent & correct multiple deficiencies



ENT18/2/02/1016



## GUARANTEED CONTENTS:

w/w

Zinc (Zn) soluble in water and chelated by EDTA:	4%
Iron (Fe) soluble in water and chelated by EDTA:	4%
Manganese (Mn) soluble in water and chelated by EDTA:	3%
Copper (Cu) soluble in water and chelated by EDTA:	0.60%
Boron (B) soluble in water:	1.50%
Sulfur (S) soluble in water:	0.96%
Magnesium (Mg) soluble in water:	0.72%
Molybdenum (Mo) soluble in water:	0.05%

EDTA pH stability range: 3-9

Distributor:  
Tradecorp APAC Pty. Ltd.  
Level 1, 225 George Street  
Sydney – NSW 2000  
Contact: 1300 595 000  
Email: [australia@tradecorp.sapec.pt](mailto:australia@tradecorp.sapec.pt)  
[www.tradecorpastralia.com.au](http://www.tradecorpastralia.com.au)

Made in Spain by:  
Trade Corporation International S.A.U.  
C/ Alcalá, 498. 2ª Planta  
28027 Madrid (España)  
Tel.: (+34) 913 273 200  
[www.tradecorp.com.es](http://www.tradecorp.com.es)  
e-mail: [global@tradecorp.com.es](mailto:global@tradecorp.com.es)

NET WEIGHT  
4 KG

# tradecorp® AZ fresco

## DOSAGE AND METHOD OF USE

Application	Crop	Rate (kg /Ha)	No. of Applications	Comments
Foliar	General Dose	0.5 - 1.5	2 - 3	Apply before or as soon as symptoms appear. 14 days apart.
	Vegetables (e.g. tomato, cucurbits)	0.5 - 1.0 (75 - 100g /100L)	2 - 3	Apply early in the crop cycle once foliage is sufficient.
	Tubers (Potato)	1	3 - 4	Apply once stems reach 10 cm. Final application post flowering. Use high volumes of water (e.g. 500L).
	Orchard Crops	1 - 1.5 (50 - 100g /100L)	2 - 4	Apply before flowering x2 - 3 and during fruit set x1.
	Cereals & Herbaceous	0.5 - 1	2 - 3	Apply at start of tillering, stem elongation and flag leaf.
	Ornamentals	0.2 - 0.5 75 - 100g /100L	2 - 3	Apply every 7 - 10 days from beginning of vegetative cycle.
Drip	General Dose	2 - 3	2 - 4	Apply every 7 - 10 days from early in the crop cycle.
Hydroponics	General Dose	3 - 5kg /1000L (x100)	-	Use 1L of mother stock solution /100L irrigation water. Use the higher rate from fruit set, high temperatures or shorter winter days.
Soil	General Dose	5 - 10	1	Apply before sowing, transplant or the beginning of plant growth.

*Spray onto dry foliage for maximum efficiency*

## DIRECTIONS FOR USE

Do not apply foliar EDTA chelates to fruit crops at >0.1% concentration (100 g /100L water). This is cumulative. If you mix two different EDTA chelate products you must not exceed 100 g /100L for the two products. i.e. maximum dosage will be 50 g of each product to give combined 100g /100L water.

Do not apply foliar EDTA chelates to broad-acre crops at >1% concentration (1 kg /100L water). The same warning about cumulative dosage applies when mixing two different EDTA products as for fruit crops above.

Do not apply by foliar application to plum or apricot trees.

Do not apply during flowering or during fruit colour change.

Add Tradecorp AZ Fresco to water while agitating and continue agitation until granules have fully dissolved.

## COMPATIBILITY

Tradecorp AZ Fresco is compatible with most commonly used fertilizers and agrochemicals. It is recommended to carry out a test before preparing any mixture and to consult technically authorized personnel. To mix with Calcium Nitrate follow these steps: 1. Fill the tank with water. 2. Add Tradecorp AZ II. 3. Mix for 2 - 3 minutes. 4. Add Calcium Nitrate. 5. Mix.

## WARNINGS, HANDLING AND STORAGE

**May damage fertility. May damage the unborn child.**

Contains: Boric acid < 10%.

Do not handle until all safety precautions have been read and understood. Use personal protective equipment as required. IF exposed or concerned: Get medical advice/attention. Dispose of contents/container according to local regulations. Keep out of reach of children. Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures. Keep only in original container. Store locked up. Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

**Disclaimer:** The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.

Manufacturing date:

Lot number:



**DANGER**