

# ultraferro

The premier Iron (Fe) EDDHA formulation to prevent & correct Iron Chlorosis, 100% water soluble



ENT18/2/02/1016



## EC FERTILIZER

### GUARANTEED CONTENTS:

w/w

Iron (Fe)	
soluble in water and chelated by EDDHA:	6.0%
Iron (Fe)	
chelated by EDDHA in ortho-ortho position:	4.2%

pH stability range: 3-11

Distributor:  
Tradecorp APAC Pty. Ltd.  
Level 1, 225 George Street  
Sydney – NSW 2000  
Contact: 1300 595 000  
Email: [australia@tradecorp.sapec.pt](mailto:australia@tradecorp.sapec.pt)  
[www.tradecorpastralia.com.au](http://www.tradecorpastralia.com.au)

Made in Spain by:  
Trade Corporation International S.A.U.  
C/ Alcalá, 498. 2ª Planta  
28027 Madrid (España)  
Tel.: (+34) 913 273 200  
[www.tradecorp.com.es](http://www.tradecorp.com.es)  
e-mail: [global@tradecorp.com.es](mailto:global@tradecorp.com.es)



NET WEIGHT  
5 KG

## DOSAGE AND METHOD OF USE

Application	Crop	Rate (kg /Ha)	No. of Applications	Comments
Fertigation	General Dose	1 - 3 1 - 2 /10,000L Water	2 - 6	Apply splits from early vegetative stage until fruit set.
Hydroponics	General Dose	1 - 1.5kg /1000L (x100)	-	Use 1L of mother stock solution /100L irrigation water.
Soil	General Dose	5 - 10	1	Apply as a single soil spray or soil injection early in the season where fertigation is not possible.
	Orchard (Establishment)	5g /tree	2	Apply at establishment.
	Orchard (Early Production)	10 - 20g /tree	2	Apply pre-flowering.
	Orchard (Full Production)	25 - 50g /tree	2	Apply pre-flowering.
	Vegetables	1 - 7g /m <sup>2</sup>	1 - 3	Apply before sowing, transplant or the beginning of plant growth.
	Flower Beds	3 - 6g /m <sup>2</sup>	1 - 3	Apply early in the season as required.
Flowers	General Dose	0.5g /L	Weekly	Water weekly.

## DIRECTIONS FOR USE

Iron EDDHA should be applied by fertigation or as a soil spray for maximum efficacy. When applied as a foliar spray Iron EDDHA is degraded by strong UV light, and is not recommended.

## COMPATIBILITY

Ultraferro is compatible with most of the fertilizers and agrochemicals. It is recommended to carry out a test before preparing any mixtures and to consult technically authorized personnel.

## WARNINGS, HANDLING AND STORAGE

Keep out of reach of children. Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures. Keep only in original container. Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians. To be used only where there is a recognised need. Do not exceed the appropriate dose rates.

**Disclaimer:** The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.



Manufacturing date:  
Lot number: