


humifirst WG

Concentrated Humic and Fulvic Acids derived from American Leonardite, the premier source of Humic Acids for agriculture, as 100% rapidly soluble micro-granule



EN18/2/02/1016



GUARANTEED CONTENTS:

w/w

Total Humic Extract (THE):	65%
Humic Acids:	53%
Fulvic Acids:	12%
Potassium (K) soluble in water:	14.11%

Distributor:
Tradecorp APAC Pty. Ltd.
Level 1, 225 George Street
Sydney – NSW 2000
Contact: 1300 595 000
Email: australia@tradecorp.sapec.pt
www.tradecorpastralia.com.au

Made in Spain by:
Trade Corporation International S.A.U.
C/ Alcalá, 498. 2ª Planta
28027 Madrid (España)
Tel.: (+34) 913 273 200
www.tradecorp.com.es
e-mail: global@tradecorp.com.es



NET WEIGHT
5 KG

humifirst WG

Humifirst WG is a natural Humic extract, based on Humic and Fulvic Acids (4:1 ratio), soluble in water and obtained from American Leonardite. Humifirst WG improves the physical, chemical and biological characteristics of soil, facilitating the availability and uptake of nutrients by crops. Humifirst WG increases the quantity of fine (white) roots. Humifirst WG is a 100% water soluble micro granule safe for use in all boom spray, sprinkler and drip irrigation systems.

Application	Crop	Rate (Kg/Ha)	No. of Applications	Comments
Drip	General (Horticulture)	1.25 - 2	2 - 5	Apply every 7 - 10 days from germination with particular focus after germination / budburst up to pre-flowering
	General (Orchard)	1.5 - 2.5	3 - 6	Apply throughout the crop cycle beginning at budburst (or pre-flowering in hysteranthus species such as almond, cherry, plum etc.). Initial applications of 2 kg /ha with a 1 - 2 post harvest applications of 1.25 kg /ha (2 kg /Ha in mechanically harvested crops like almonds).
Flood	General	1.75 - 2.5	4 - 6	Apply 1 week after emergence. Subsequent applications should target maximum growth stages, stem elongation, flowering and fruit set.
	Cane	3.5 - 5	1	Apply at each ratoon
Sprinkler / Pivot	General	1.75 - 2.5	3 - 5	Apply as per drip irrigation
Soil Spray (Boom)	General	7.5 - 10	1	Apply early in the crop cycle, pre, during or post sowing. Use a minimum of 200 L /Ha water
Soil Spray (Row)	General	4 - 5	1	Apply early in the cycle, pre, during or post sowing. Use a minimum of 150 L /Ha water
Growth Substrate	General	400 g /m ³	1	Incorporate when preparing the potting mix
	Sprinkler	2.5 g /L	1 - 2	In situations of young foliage contact apply at 0.25% concentration. Do not apply when leaf cover of soil exceeds 25%


COMPATIBILITY

Humifirst WG is compatible with most commonly used fertilizers and agrochemicals, except mineral oils, calcium nitrate, acid reaction products and herbicides incompatible with organic matter. It is recommended to carry out a test before preparing any mixture and to consult technically authorized personnel.

HANDLING AND STORAGE

Keep out of the reach of children. Do not eat, drink or smoke when using this product. Store in a fresh, dry place and keep away from extreme temperatures. Keep only in original container. Use the product according to the instructions in this label. In case of doubt, please contact authorized technicians. To optimise the speed of dissolution and to avoid the formation of lumps, it is recommended that the product be added slowly to the mixing tank with the agitator running and mixed until total dissolution is achieved.

Disclaimer: The recommendations and information provided comes from several trials and studies that have been undertaken, however, during the application of the product several factors out of our control may take place (tank mixes, application, climatic conditions, etc.). Any recommendation must serve as a guide and be adapted to local conditions.



Manufacturing date:
Lot number:

8 435146 607387